

ABSOLUTELY SUMMING OPERATORS ON NON COMMUTATIVE C^* -ALGEBRAS AND APPLICATIONS

NARCISSE RANDRIANANTOANINA

ABSTRACT. Let E be a Banach space that does not contain any copy of ℓ^1 and $\mathcal{A}$ be a non commutative C^* -algebra. We prove that every absolutely summing operator from $\mathcal{A}$ into E^* is compact, thus answering a question of Pełczyński.

As application, we show that if G is a compact metrizable abelian group and Λ is a Riesz subset of its dual then every countably additive $\mathcal{A}^*$ -valued measure with bounded variation and whose Fourier transform is supported by Λ has relatively compact range. Extensions of the same result to symmetric spaces of measurable operators are also presented.

1. INTRODUCTION

It is a well known result that every absolutely summing operator from a $C(K)$ -space into a separable dual space is compact. More generally if F is a Banach space with the complete continuity property (CCP) then every absolutely summing operator from any $C(K)$ -spaces into F is compact (see [10]).

It is the intention of the present note to study extensions of the above results in the setting of C^* -algebras, i.e., replacing the $C(K)$ -spaces above by a general non commutative C^* - algebra. Typical examples of Banach spaces with the CCP are dual spaces whose preduals do not contain ℓ^1 . Our main result is that if E is a Banach space that does not contain any copy of ℓ^1 and $\mathcal{A}$ is a C^* -algebra then every absolutely summing operator from $\mathcal{A}$ into E^* is compact. This answered positively the following question raised by Pełczyński (see [17] Problem 3. P.20): Is every absolutely summing operator from a non commutative C^* -algebra into a Hilbert space compact? This result is also used to study

1991 *Mathematics Subject Classification.* 46E40; Secondary 47D15, 28B05.

Key words and phrases. C^* -algebras, vector measures, Riesz sets.

relative compactness of range of countably additive vector measures with values in duals of non commutative C^* -algebras. In [9], Edgar introduced new types of Radon-Nikodym properties associated with Riesz subsets of countable discrete group (see the definition below) as generalization of the usual Radon-Nikodym property (RNP) and the Analytic Radon-Nikodym property (ARNP). These properties were extensively studied in [7] and [8]. In [7], it was shown that if Λ is a Riesz subset of a countable discrete group then $L^1[0, 1]$ has the type II- Λ -RNP. In the other hand, Haagerup and Pisier showed in [14] that non commutative L^1 -spaces have the ARNP so it is a natural question to ask if non commutative L^1 -spaces have the type II- Λ -RNP for any Riesz subset. In this direction, we obtain (as a consequence of our main result) that if a countably additive vector measure of bounded variation is defined on the σ -field of Borel subsets of a compact metrizable abelian group and takes its values in a dual of a C^* -algebra then its range is relatively compact provided that its Fourier transform is supported by a Riesz subset of the dual group.

Our terminology and notation are standard. We refer to [3] and [4] for definitions from Banach space theory and [21], [16] and [11] for basic properties from the theory of operator algebras and non-commutative integrations.

2. PRELIMINARY FACTS AND NOTATIONS

We recall some definitions and well known facts which we use in the sequel.

Let $\mathcal{A}$ be a C^* -algebra, we denote by $\mathcal{A}_h$ the set of Hermitian (self adjoint) elements of $\mathcal{A}$.

Definition 1. *Let E and F be Banach spaces and $0 < p < \infty$. An operator $T : E \rightarrow F$ is said to be absolutely p -summing (or simply p -summing) if there exists C such that*

$$\left(\sum_{i=1}^{\ell} \|Te_i\|^p \right)^{1/p} \leq C \max \left\{ \sum |\langle e_i, e^* \rangle|^p, \|e^*\| \leq 1 \right\}^{1/p}$$

The following class of operators was introduced by Pisier in [18] as extension of the q -summing operators in the setting of C^* -algebras.

Definition 2. *Let $\mathcal{A}$ be a C^* -algebra and F be a Banach space, $0 < q < \infty$. An operator $T : \mathcal{A} \rightarrow F$ is said to be q - C^* -summing if there exists a constant C such that for any finite sequence $(A_1, \dots, A_n)$ of Hermitian elements of $\mathcal{A}$ one has*

$$\left(\sum_{i=1}^n \|T(A_i)\|^q \right)^{1/q} \leq C \left\| \left(\sum_{i=1}^n |A_i|^q \right)^{1/q} \right\|_{\mathcal{A}}.$$

The smallest constant C for which the above inequality holds is denoted by $C_q(T)$. It should be noted that if the C^* -algebra $\mathcal{A}$ is commutative then every q - C^* -summing operator from $\mathcal{A}$ into any Banach space is q -summing. The following extension of the classical Pietsch's factorization theorem ([3]) was obtained by Pisier (see Proposition 1.1 of [18]).

Proposition 1. *If $T : \mathcal{A} \rightarrow F$ is a q - C^* -summing operator then there exists a positive linear form f of norm less than 1 such that*

$$\|Tx\| \leq C_q(T) \{f(|x|^q)\}^{1/q}, \text{ for every } x \in \mathcal{A}_h.$$

Let $\mathcal{M}$ be a von-Neumann algebra and $\mathcal{M}_*$ be its predual. We recall that a functional f on $\mathcal{M}$ is called normal if it belongs to $\mathcal{M}_*$. In [19], it was shown that for the case of von-Neumann algebra and the operator T being weak* to weakly continuous then the positive linear form on the above proposition can be chosen to be normal; namely we have the following lemma (see Lemma 4.1 of [19]).

Lemma 1. *Let $T : \mathcal{M} \rightarrow F$ be a 1- C^* -summing operator. If T is weak* to weakly continuous then there exists a linear form $f \in \mathcal{M}_*$ with $\|f\| \leq 1$ such that*

$$\|Tx\| \leq C_1(T) f(|x|), \text{ for every } x \in \mathcal{M}_h. \quad (1)$$

For the next lemma, we recall that for $x \in \mathcal{M}$ and $f \in \mathcal{M}^*$, xf (resp. fx) denotes the element of $\mathcal{M}^*$ defined by $xf(y) = f(yx)$ (resp. $fx(y) = f(xy)$) for all $y \in \mathcal{M}$.

Lemma 2. *Let f be a positive linear form on $\mathcal{M}$. For every $x \in \mathcal{M}$,*

$$f\left(\left(\frac{xx^* + x^*x}{2}\right)^{1/2}\right) \leq 2\|xf + fx\|_{\mathcal{M}^*}. \quad (2)$$

Proof. Assume first that $x \in \mathcal{M}_h$. In this case $(xx^* + x^*x)/2 = |x|^2$. The operator x can be decomposed as $x = x_+ - x_-$ where $x_+, x_- \in \mathcal{M}^+$ and $x_+x_- = 0$. There exists a projection $p \in \mathcal{M}$ such that $px_- = x_-p = x_-$ and $(1-p)x_+ = x_+(1-p) = x_+$. This yields the following estimates:

$$\begin{aligned} f(|x|) &= f(x_+ + x_-) = f(x_+) + f(x_-) \\ &= \frac{1}{2}(xf + fx)(1-p) + \frac{1}{2}(xf + fx)(p) \\ &\leq \frac{1}{2}(\|xf + fx\|\|1-p\| + \|xf + fx\|\|p\|) \\ &\leq \|xf + fx\|. \end{aligned}$$

For the general case, fix $x \in \mathcal{M}$. Let $a = (x + x^*)/2$ and $b = (x - x^*)/2i$. Clearly $x = a + ib$ and $((xx^* + x^*x)/2)^{1/2} = |a| + |b|$. Using the Hermitian case, we get:

$$\begin{aligned} f(|a| + |b|) &\leq \|af + fa\| + \|bf + fb\| \\ &\leq \|xf + fx\| + \|x^*f + fx^*\|; \end{aligned}$$

but since $f \geq 0$,

$$\begin{aligned} \|x^*f + fx^*\| &= \sup\{|f(sx^* + x^*s)|; s \in \mathcal{M}, \|s\| \leq 1\} \\ &= \sup\{|f^*(xs^* + s^*x)|; s \in \mathcal{M}, \|s\| \leq 1\} \\ &= \sup\{|f(xs^* + s^*x)|; s \in \mathcal{M}, \|s\| \leq 1\} \\ &= \|xf + fx\|, \end{aligned}$$

which completes the proof of the lemma. $\square$

3. MAIN THEOREM

Theorem 1. *Let $\mathcal{A}$ be a C^* -algebra, E be a Banach space that does not contain any copy of ℓ^1 and $T : \mathcal{A} \rightarrow E^* = F$ be a 1-summing operator then T is compact.*

We will divide the proof into two steps. First we will assume that the C^* -Algebra $\mathcal{A}$ is a σ -finite von-Neumann algebra and the operator T is weak* to weakly continuous; then we will show that the general can be reduced to this case. We refer to [21] P. 78 for the definition of σ -finite von-Neumann algebra.

Proposition 2. *Let $\mathcal{M}$ be a σ -finite von-Neumann algebra. Let $T : \mathcal{M} \rightarrow E^*$ be a weak* to weakly continuous 1-summing operator then T is compact.*

Proof. The operator T being weak* to weakly continuous and 1-summing, there exist a constant $C = C_1(T)$ and a normal positive functional f on $\mathcal{M}$ such that

$$\|Tx\| \leq Cf(|x|) \text{ for every } x \in \mathcal{M}_h.$$

Since the von-Neumann algebra $\mathcal{M}$ is σ -finite, there exists a faithful normal state f_0 in $\mathcal{M}_*$ (see [21] Proposition II-3.19). Replacing f by $f + f_0$, we can assume that the functional f on the inequality above is a faithful normal state and using Lemma 2, we get

$$\|Tx\| \leq 2C\|xf + fx\|_{\mathcal{M}^*} \text{ for every } x \in \mathcal{M}. \quad (3)$$

We may equip $\mathcal{M}$ with the scalar product by setting for every $x, y \in \mathcal{M}$,

$$\langle x, y \rangle = f\left(\frac{xy^* + y^*x}{2}\right).$$

Since f is faithful, $\mathcal{M}$ with $\langle \cdot, \cdot \rangle$ is a pre-Hilbertian. we denote the completion of this space by $L^2(\mathcal{M}, f)$ (or simply $L^2(f)$).

By construction, the inclusion map $J : \mathcal{M} \rightarrow L^2(\mathcal{M}, f)$ is bounded and is one to one (f is faithful). On the dense subspace $J(\mathcal{M})$ of $L^2(f)$, we define a map $\theta : J(\mathcal{M}) \rightarrow L^2(f)^*$ by $\theta(Jx) = \langle \cdot, J(x^*) \rangle$. The map θ is clearly linear and is an isometry; indeed for every $x \in \mathcal{M}$, $\|\theta(Jx)\|^2 = \sup_{\|u\| \leq 1} \langle u, J(x^*) \rangle^2 = \langle J(x^*), J(x^*) \rangle = f(x^*x + xx^*) = \|Jx\|^2$.

So it can be extended to a bounded map (that we will denote also by θ) from $L^2(f)$ onto $L^2(f)^*$.

Let $S = J^* \circ \theta \circ J$. The operator is defined from $\mathcal{M}$ into $\mathcal{M}_*$ and we claim that for every $x \in \mathcal{M}$, $Sx = xf + fx$. In fact for every $x, y \in \mathcal{M}$, we have:

$$\begin{aligned} Sx(y) &= J^* \circ \theta \circ Jx(y) \\ &= \theta \circ Jx(Jy) \\ &= \langle J(y), J(x^*) \rangle \\ &= f(xy + yx) = (xf + fx)(y). \end{aligned}$$

Notice also that since f is normal, the functionals xf and fx are both normal for every $x \in \mathcal{M}$; therefore $S(\mathcal{M}) \subset \mathcal{M}_*$. Also since J is one to one, J^* has weak* dense range. The latter with the facts that both J and θ have dense ranges imply that $S(\mathcal{M})$ is weak* dense in $\mathcal{M}^*$ so $S(\mathcal{M})$ is (norm) dense in $\mathcal{M}_*$.

Let us now define a map $L : S(\mathcal{M}) \rightarrow E^*$ by $L(xf + fx) = Tx$ for every $x \in \mathcal{M}$. The map L is clearly linear and one can deduce from (3) that L is bounded so it can be extended as a bounded operator (that we will denote also by L) from $\mathcal{M}_*$ into E^* . The above means that T can be factored as follows

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{T} & E^* \\ S \searrow & & \nearrow L \\ & \mathcal{M}_* & \end{array}$$

Taking the adjoints we get

$$\begin{array}{ccc} E & \xrightarrow{T^*} & \mathcal{M}_* \\ L^* \searrow & & \nearrow S^* \\ & \mathcal{M} & \end{array}$$

To conclude the proof of the proposition, let $(e_n)_n$ be a bounded sequence in E . Since $E \not\hookrightarrow \ell^1$, we will assume (by taking a subsequence if necessary) that $(e_n)_n$ is weakly Cauchy. We will show that $(T^*(e_n))_n$ is norm-convergent. For that it is enough to prove that if $(e_n)_n$ is a weakly null sequence in E then $(\|T^*e_n\|)_n$ converges to zero.

Let $(e_n)_n$ be a weakly null sequence in E , $(L^*(e_n))_n$ is a weakly null sequence in $\mathcal{M}$. This implies that $((L^*(e_n))^*)_{n \geq 1}$ (the sequence of the adjoints of the $L^*(e_n)$'s) is weakly null in $\mathcal{M}$.

Since T is 1-summing, it is a Dunford-Pettis operator (i.e takes weakly convergent sequence into norm-convergent sequence). Hence

$$\lim_{n \rightarrow \infty} \|T((L^*e_n)^*)\|_{E^*} = 0.$$

In particular, since $(e_n)_n$ is a bounded sequence in E , we have

$$\lim_{n \rightarrow \infty} \langle T((L^*e_n)^*), e_n \rangle = 0$$

but

$$\begin{aligned} \langle T((L^*e_n)^*), e_n \rangle &= \langle LS((L^*e_n)^*), e_n \rangle \\ &= \langle S((L^*e_n)^*), L^*e_n \rangle \\ &= \langle \theta \circ J((L^*e_n)^*), J(L^*e_n) \rangle \\ &= \langle J(L^*e_n), J(L^*e_n) \rangle_{L^2(f)} \\ &= \|J(L^*e_n)\|_{L^2(f)}. \end{aligned}$$

So $\|J(L^*e_n)\|_{L^2(f)} \rightarrow 0$ as $n \rightarrow \infty$ and therefore since $T^* = S^* \circ L^* = J^* \circ \theta \circ J \circ L^*$, we get that $\lim_{n \rightarrow \infty} \|T^*e_n\| = 0$.

This shows that $\overline{T^*(B_E)}$ is compact and since B_E is weak* dense in $B_{E^{**}}$ and T^* is weak* to weakly continuous, $T^*(B_{E^{**}}) \subseteq \overline{T^*(B_E)}$ so T^* (and hence T) is compact. The proposition is proved. $\square$

To complete the proof of the theorem, let $\mathcal{A}$ be a C^* -algebra and $T : \mathcal{A} \rightarrow E^*$ be a 1-summing operator. The double dual $\mathcal{A}^{**}$ of $\mathcal{A}$ is a von-Neumann and $T^{**} : \mathcal{A}^{**} \rightarrow E^*$ is 1-summing. Let $(a_n)_n$ be a bounded sequence in $\mathcal{A}^{**}$. If we denote by $\mathcal{M}$ the von-Neumann algebra generated by $(a_n)_n$ then the predual $\mathcal{M}_*$ of $\mathcal{M}$ is separable and therefore the von-Neumann algebra $\mathcal{M}$ is σ -finite. Moreover, if we set $I : \mathcal{M} \rightarrow \mathcal{A}^{**}$ the inclusion map then I is weak* to weak* continuous. Hence $\mathcal{M}$ and $T^{**} \circ I$ satisfy the conditions of Proposition 2 so $T^{**} \circ I$ is compact and since the sequence $(a_n)_n$ is arbitrary, the operator T^{**} (and hence T) is compact. $\square$

Remark. It should be noted that for the proof of Proposition 2, we only require the operator T to be C^* -summing and Dunford-Pettis so

the conclusion of Proposition 2 is still valid for C^* -summing operators that are Dunford-Pettis.

4. APPLICATIONS TO VECTOR MEASURES

In this section we will provide some applications of the main theorem to study range of countably additive vector measures with values in duals of C^* -Algebras.

The letter G will denote a compact metrizable abelian group, $\widehat{G}$ its dual, $\mathcal{B}(\mathcal{G})$ is the σ -algebra of the Borel subsets of G , and λ the normalized Haar measure on G .

Let X be a Banach space and $1 \leq p \leq \infty$, we will denote by $L^p(G, X)$ the usual Bochner spaces for the measure space $(G, \mathcal{B}(\mathcal{G}), \lambda)$; $M(G, X)$ the space of X -valued countably additive Borel measures of bounded variation; $C(G, X)$ the space of X -valued continuous functions and $M^\infty(G, X) = \{\mu \in M(G, X), |\mu| \leq C\lambda \text{ for some } C > 0\}$.

If $\mu \in M(G, X)$, we recall that the Fourier transform of μ is a map $\hat{\mu}$ from $\widehat{G}$ into X defined by $\hat{\mu}(\gamma) = \int_G \bar{\gamma} d\mu$ for $\gamma \in \widehat{G}$.

For $\Lambda \subset \widehat{G}$, we will use the following notation:

$$\begin{aligned} L_\Lambda^p(G, X) &= \{f \in L^p(G, X), \hat{f}(\gamma) = 0 \text{ for all } \gamma \notin \Lambda\} \\ C_\Lambda(G, X) &= \{f \in C(G, X), \hat{f}(\gamma) = 0 \text{ for all } \gamma \notin \Lambda\} \\ M_\Lambda(G, X) &= \{\mu \in M(G, X), \hat{\mu}(\gamma) = 0 \text{ for all } \gamma \notin \Lambda\} \\ M_\Lambda^\infty(G, X) &= \{\mu \in M^\infty(G, X), \hat{\mu}(\gamma) = 0 \text{ for all } \gamma \notin \Lambda\}. \end{aligned}$$

We also recall that $\Lambda \subset \widehat{G}$ is called a Riesz subset if $M_\Lambda(G) = L_\Lambda^1(G)$. We refer to [20] and [13] for detailed discussions and examples of Riesz subsets of dual groups.

The following Banach space properties were introduced by Edgar [9], and Dowling [7].

Definition 3. *Let Λ be a Riesz subset of $\widehat{G}$. A Banach space X is said to have type I- Λ -Radon Nikodym Property (resp. type II- Λ -Radon Nikodym property) if $M_\Lambda^\infty(G, X) = L_\Lambda^\infty(G, X)$ (resp. $M_\Lambda(G, X) = L_\Lambda^1(G, X)$).*

Our next result deals with property of dual of C^* -algebras related to the types of Radon-Nikodym properties defined above.

Theorem 2. *Let Λ be a Riesz subset of $\widehat{G}$ and $\mathcal{A}$ be a C^* -Algebra. If $F : \mathcal{B}(G) \rightarrow \mathcal{A}^*$ is a countably additive measure with bounded variation that satisfies $\widehat{F}(\gamma) = 0$ for $\gamma \notin \Lambda$ then the range of F is a relatively compact subset of $\mathcal{A}^*$.*

Proof. Let $F : \mathcal{B}(G) \rightarrow \mathcal{A}^*$ be a measure with bounded variation and $\widehat{F}(\gamma) = 0$ for $\gamma \notin \Lambda$. Let $S : C(G) \rightarrow \mathcal{A}^*$ be the operator defined by $Sf = \int f dF$. Since F is of bounded variation, the operator S is integral (see [4] Theorem IV-3.3 and Theorem IV-3.12) and therefore $S^* : \mathcal{A}^{**} \rightarrow (C(G))^*$ is also integral. Now since $\widehat{F}(\gamma) = 0$ for $\gamma \notin \Lambda$, if we denote by $\Lambda' = \{\gamma \in \widehat{G}, \bar{\gamma} \notin \Lambda\}$ then $S(\gamma) = 0$ for all $\gamma \in \Lambda'$ and therefore we have the following factorization

$$\begin{array}{ccc} C(G) & \xrightarrow{S} & \mathcal{A}^* \\ q \downarrow & \nearrow_L & \\ C(G)/C_{\Lambda'}(G) & & \end{array}$$

where q is the natural quotient map. Taking the adjoints, we get

$$\begin{array}{ccc} \mathcal{A}^{**} & \xrightarrow{S^*} & (C(G))^* \\ L^* \searrow & & \nearrow_{q^*} \\ & M_\Lambda(G) & \end{array}$$

Since q^* is the formal inclusion and S^* is 1-summing, the operator L^* is 1-summing. The assumption Λ being a Riesz subset implies that $M_\Lambda(G) = L_\Lambda^1(G)$ is a separable dual (in particular its predual does not contain ℓ^1). So by Theorem 1, L^* (and hence S) is compact. This proves that the range of the representing measure F of S is relatively compact (see [4] Theorem II-2.18). $\square$

Our next result is a generalization of Theorem 2 for the case of symmetric spaces of measurable operators.

Let $(\mathcal{M}, \tau)$ be a semifinite von-Neumann algebra acting on a Hilbert space H . Let τ be a distinguished faithful normal semifinite trace on $\mathcal{M}$.

Let $\overline{\mathcal{M}}$ be the space of all measurable operators with respect to $(\mathcal{M}, \tau)$ in the sense of [16]; for $a \in \overline{\mathcal{M}}$ and $t > 0$, the t^{th} -s-number (singular number) of a is defined by

$$\mu_t(a) = \inf\{\|ae\| : e \in \mathcal{M} \text{ projection with } \tau(I - e) \leq t\}.$$

The function $t \mapsto \mu_t(a)$ defined on $(0, \tau(I))$ will be denoted by $\mu(a)$. This is a positive non-increasing function on $(0, \tau(I))$. We refer to [11] for complete detailed study of $\mu(a)$.

Let E be a rearrangement invariant Banach function space on $(0, \tau(I))$ (in the sense of [15]). We define the symmetric space $E(\mathcal{M}, \tau)$ of measurable operators by setting

$$E(\mathcal{M}, \tau) = \{a \in \overline{\mathcal{M}}; \mu(a) \in E\}$$

and $\|a\|_{E(\mathcal{M}, \tau)} = \|\mu(a)\|_E$.

It is well known that $E(\mathcal{M}, \tau)$ is a Banach space and if $E = L^p(0, \tau(I))$ ($1 \leq p \leq \infty$) then $E(\mathcal{M}, \tau)$ coincide with the usual non-commutative L^p -spaces associated with the von-Neumann algebra $\mathcal{M}$. The space $E(\mathcal{M}, \tau)$ is often referred as the non-commutative version of the function space E . Some Banach space properties of these spaces can be found in [2], [6] and [22].

For the case where the trace τ is finite, we obtain the following generalization of Theorem 2 for symmetric spaces of measurable operators.

Corollary 1. *Assume that τ is finite. Let E be a rearrangement invariant function space on $(0, \tau(I))$ that does not contain c_0 and Λ be a Riesz subset of $\widehat{G}$. Let $F : \mathcal{B}(G) \rightarrow E(\mathcal{M}, \tau)$ be a countably additive measure with bounded variation and $\widehat{F}(\gamma) = 0$ for every $\gamma \notin \Lambda$ then the range of F is relatively compact.*

Proof. We will begin by reducing the general case to the case where $E(\mathcal{M}, \tau)$ is separable. Since $\mathcal{B}(G)$ is countably generated, the range of F is separable. Choose $(A_n)_n \subset \mathcal{B}(G)$ so that $\{F(A_n), n \geq 1\}$ is dense in $\{F(A), A \in \mathcal{B}(G)\}$. Let $\widetilde{\mathcal{M}}$ be the von-Neumann algebra generated I and $F(A_n)$ ($n \geq 1$) and $\widetilde{\tau}$ the restriction of τ in $\widetilde{\mathcal{M}}$. Clearly $E(\widetilde{\mathcal{M}}, \widetilde{\tau})$ is a closed subspace of $E(\mathcal{M}, \tau)$ and $F(A) \in E(\widetilde{\mathcal{M}}, \widetilde{\tau})$ for all $A \in \mathcal{B}(G)$.

Moreover the space $E(\widetilde{\mathcal{M}}, \tilde{\tau})$ is separable (see Lemma 5.6 of [22]). So without loss of generalities we will assume that $E(\mathcal{M}, \tau)$ is separable. It is a well known fact that $E(\mathcal{M}, \tau)$ is contained in $L^1(\mathcal{M}, \tau) + \mathcal{M}$ and since τ is finite, $E(\mathcal{M}, \tau) \subset L^1(\mathcal{M}, \tau)$. Let $J : E(\mathcal{M}, \tau) \rightarrow L^1(\mathcal{M}, \tau)$ be the formal inclusion. The measure $J \circ F$ is of bounded variation and $\widehat{J \circ F}(\gamma) = J(\widehat{F}(\gamma))$ for every $\gamma \in \widehat{G}$. One can conclude from Theorem 2 that the range of $J \circ F$ is relatively compact in $L^1(\mathcal{M}, \tau)$.

To show that the range of F is relatively compact, fix $h : G \rightarrow E(\mathcal{M}, \tau)^{**}$ a weak*-density of F with respect to the Haar measure λ (see [5]). We have for each $A \in \mathcal{B}(G)$,

$$F(A) = \text{weak}^* - \int_A h(t) d\lambda(t)$$

and

$$|F|(A) = \int_A \|h(t)\| d\lambda(t).$$

For each $N \in \mathbb{N}$, let $A_N = \{t \in G, \|h(t)\| \leq N\}$ and F_N the measure defined by $F_N(A) = F(A \cap A_N)$ for all $A \in \mathcal{B}(G)$. Clearly $|F_N| \leq N\lambda$ for every $N \in \mathbb{N}$.

Define $T_N : L^1(G) \rightarrow E(\mathcal{M}, \tau)$ by $T_N(f) = \int f(t) dF_N(t)$ for every $f \in L^1(G)$. The operator T_N is bounded and we claim that T_N is Dunford-Pettis; for that notice that since the range of $J \circ F$ is relatively compact so is the range of $J \circ F_N$ and therefore the operator $J \circ T_N$ is a Dunford-Pettis operator. The space $E(\mathcal{M}, \tau)$ is separable and J is a semi-embedding (see Lemma 5.7 of [22]) so J is a G_δ -embedding (see [1] Proposition 1.8) and one can deduce from Theorem II.6 of [12], that T_N is a Dunford-Pettis operator. Hence the range of F_N is relatively compact. Now since

$$\lim_{N \rightarrow \infty} \|F - F_N\| = \lim_{N \rightarrow \infty} \int_{G \setminus A_N} \|h(t)\| d\lambda(t) = 0,$$

the range of F is relatively compact. $\square$

Let us finish by asking the following question:

Question: Do non-commutative L^1 -spaces have type II- Λ -RNP for any Riesz set Λ ?

In light of Theorem 2, the result of Haagerup and Pisier ([14]) and so many properties that have been generalized from classical L^1 -spaces to non-commutative L^1 -spaces, one tends to conjecture that the answer of the above question is affirmative.

Acknowledgements: I would like to thank Professor G. Pisier for many valuable suggestions concerning this work.

REFERENCES

- [1] J. Bourgain and H. P. Rosenthal. *Applications of the theory of semi-embeddings to Banach space theory*. J. Funct. Anal., **52**:149–188, (1983).
- [2] V.I. Chilin and F.A. Sukochev. *Symmetric spaces on semifinite von Neumann algebras*. Soviet Math. Dokl., **42**:97–101, (1992).
- [3] J. Diestel. *Sequences and Series in Banach Spaces*, volume **92** of Graduate Text in Mathematics. Springer Verlag, New York, first edition, (1984).
- [4] J. Diestel and Jr. J.J. Uhl. *Vector Measures*, volume **15** of Math Surveys. AMS, Providence, RI, (1977).
- [5] N. Dinculeanu. *Vector Measures*. Pergamon Press, New York, (1967).
- [6] P.G. Dodds, T.K. Dodds, and B. Pagter. *Non-commutative Banach function spaces*. Math. Zeit., **201**:583–597, (1989).
- [7] P. Dowling. *Radon-Nikodym properties associated with subsets of countable discrete abelian groups*. Trans. Amer. Math. Soc., **327**:879–890, (1991).
- [8] P. Dowling. *Duality in some vector-valued function spaces*. Rocky Mount. Jr. of Math., **22**:511–518, (1992).
- [9] G. Edgar. *Banach spaces with the analytic Radon-Nikodym property and compact abelian group*. Proc. Int. Conf. on Almost Everywhere convergence in Probability and Ergodic Theory, Academic Press, pages 195–213, (1989).
- [10] G. Emmanuele. *Dominated operators on $C[0, 1]$ and the (CRP)*. Collect. Math., **41**:21–25, (1990).
- [11] T. Fack and H. Kosaki. *Generalized s -numbers of τ -measurable operators*. Pac. J. Math., **123**:269–300, (1986).
- [12] N. Ghoussoub and H. P. Rosenthal. *Martingales, G_δ -Embeddings and quotients of L_1* . Math. Ann., **264**:321–332, (1983).
- [13] G. Godefroy. *On Riesz subsets of Abelian discrete groups*. Isr. J. of Math., **61**:301–331, (1988).
- [14] U. Haagerup and G. Pisier. *Factorization of Analytic functions with values in non-commutative L_1 -spaces*. Canad. J. Math., **41**:882–906, (1989).
- [15] J. Lindenstrauss and L. Tzafriri. *Classical Banach spaces II*, volume **97** of Modern Survey In Mathematics. Springer-Verlag, Berlin-Heidelberg-New York, first edition, (1979).
- [16] E. Nelson. *Notes on non-commutative integration*. J. Funct. Anal., **15**:103–116, (1974).

- [17] A. Pelczynski. *Compactness of absolutely summing operators*. In V. Havin and N. Nikolski, editors, *Linear and Complex Analysis*, volume **1573** of Lectures Notes in Mathematics, Berlin Heidelberg New York, (1994). Springer Verlag.
- [18] G. Pisier. *Grothendieck's Theorem for non-commutative C^* -algebras with appendix on Grothendieck's constants*. J. Func. Anal., **29**:397–415, (1978).
- [19] G. Pisier. *Factorization of operators through $L_{p\infty}$ or $L_{1\infty}$ and Non-commutative generalizations*. Math. Ann., **276**:105–136, (1986).
- [20] W. Rudin. *Fourier analysis on groups*, volume **12** of Interscience Tracts in Pure and Appl. Math. . Interscience, New York, first edition, (1962).
- [21] M. Takesaki. *Theory of operator Algebras I*. Springer-Verlag, New-York, Heidelberg, Berlin, (1979).
- [22] Q. XU. *Analytic functions with values in Lattices and Symmetric spaces of measurable operators*. Math. Proc. Camb. Phil. Soc., **109**:541–563, (1991).

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN,
AUSTIN, TX 78712-1082

E-mail address: nrandri@math.utexas.edu